

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>All causes</b>															
2000 .....	872.0	1,042.5	739.1	585.6	718.8	475.7	886.1	1,058.3	752.7	861.9	1,028.6	732.6	1,161.1	1,417.2	973.5
1999 .....	881.9	1,061.8	743.6	601.0	736.0	490.1	894.9	1,076.6	756.0	869.4	1,045.1	734.9	1,177.0	1,451.0	979.3
<b>Salmonella infections (A01-A02)</b>															
2000 .....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
1999 .....	0.0	0.0	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
<b>Shigellosis and amebiasis (A03,A06)</b>															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Certain other intestinal infections (A04,A07-A09)</b>															
2000 .....	0.5	0.5	0.5	0.3	0.3	0.2	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.3
1999 .....	0.4	0.4	0.4	0.2	*	*	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.4
<b>Tuberculosis (A16-A19)</b>															
2000 .....	0.3	0.4	0.2	0.5	0.6	0.3	0.2	0.4	0.2	0.1	0.2	0.1	1.0	1.5	0.6
1999 .....	0.3	0.5	0.2	0.7	1.0	0.5	0.3	0.5	0.2	0.2	0.3	0.1	1.0	1.5	0.6
<b>Respiratory tuberculosis (A16)</b>															
2000 .....	0.2	0.3	0.1	0.3	0.6	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.8	1.3	0.4
1999 .....	0.3	0.4	0.2	0.6	0.7	0.4	0.3	0.4	0.2	0.2	0.3	0.1	0.8	1.2	0.5
<b>Other tuberculosis (A17-A19)</b>															
2000 .....	0.1	0.1	0.0	0.1	*	*	0.0	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.2
1999 .....	0.1	0.1	0.0	0.1	0.2	*	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.3	*
<b>Whooping cough (A37)</b>															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Scarlet fever and erysipelas (A38,A46)</b>															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Meningococcal infection (A39)</b>															
2000 .....	0.0	0.1	0.1	0.1	*	*	0.0	0.1	0.1	0.0	0.0	0.1	0.1	0.2	*
1999 .....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
<b>Septicemia (A40-A41)</b>															
2000 .....	11.4	12.2	10.6	7.0	8.1	6.2	11.5	12.4	10.8	10.2	11.1	9.6	25.4	28.0	23.5
1999 .....	11.3	12.4	10.5	7.7	8.7	6.8	11.5	12.6	10.7	10.2	11.1	9.5	25.5	28.9	23.0
<b>Syphilis (A50-A53)</b>															
2000 .....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
1999 .....	0.0	*	0.0	*	*	*	0.0	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute poliomyelitis (A80)															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)															
2000 .....	1.9	2.6	1.3	3.4	4.4	2.4	1.8	2.5	1.2	1.6	2.2	1.1	2.8	4.0	1.9
1999 .....	1.8	2.4	1.2	3.4	4.2	2.6	1.6	2.3	1.1	1.5	2.0	1.0	2.7	3.6	1.9
Human immunodeficiency virus (HIV) disease (B20-B24)															
2000 .....	5.3	8.0	2.6	7.0	11.1	3.0	5.0	7.6	2.5	2.3	3.8	0.7	24.4	37.1	13.9
1999 .....	5.4	8.4	2.6	7.3	11.5	3.1	5.2	8.0	2.5	2.4	4.1	0.7	24.9	38.2	13.8
Malaria (B50-B54)															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)															
2000 .....	2.0	2.4	1.7	1.9	2.3	1.6	2.0	2.4	1.7	1.9	2.2	1.6	3.1	3.8	2.6
1999 .....	2.7	3.1	2.3	2.4	2.6	2.3	2.7	3.1	2.4	2.4	2.8	2.1	4.8	5.6	4.2
Malignant neoplasms (C00-C97)															
2000 .....	201.0	247.4	169.6	120.8	150.1	100.6	205.6	253.0	173.7	202.6	247.5	172.5	256.8	342.9	202.3
1999 .....	202.7	251.6	169.9	122.0	151.4	101.4	207.2	257.1	173.8	203.8	251.4	172.1	261.4	350.1	205.6
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)															
2000 .....	2.7	4.1	1.6	1.6	2.8	0.7	2.8	4.2	1.7	2.6	3.8	1.6	4.2	7.5	1.8
1999 .....	2.8	4.2	1.6	1.7	2.8	0.9	2.8	4.3	1.7	2.6	3.9	1.6	4.5	7.8	2.2
Malignant neoplasm of esophagus (C15)															
2000 .....	4.4	7.7	1.8	2.3	4.0	1.0	4.6	8.0	1.9	4.4	7.8	1.8	6.7	11.3	3.4
1999 .....	4.4	7.7	1.8	2.3	4.2	0.9	4.5	7.9	1.9	4.4	7.7	1.7	6.7	11.8	3.2

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of stomach (C16)															
2000 .....	4.6	6.4	3.3	6.3	8.7	4.6	4.4	6.2	3.2	3.8	5.3	2.6	9.3	12.8	6.9
1999 .....	4.7	6.6	3.3	6.4	8.4	4.8	4.5	6.4	3.2	3.9	5.5	2.7	9.5	13.7	6.7
Malignant neoplasms of colon, rectum and anus (C18-C21)															
2000 .....	20.9	24.9	17.9	12.4	15.5	10.2	21.3	25.4	18.3	20.8	24.7	17.7	29.2	35.3	25.1
1999 .....	21.1	25.2	18.1	12.8	15.6	10.7	21.5	25.7	18.4	21.0	25.2	17.8	29.3	34.5	25.8
Malignant neoplasms of liver and intrahepatic bile ducts (C22)															
2000 .....	4.7	6.8	3.0	6.7	9.7	4.4	4.5	6.6	2.9	4.0	5.8	2.6	6.3	9.8	3.8
1999 .....	4.6	6.5	3.0	6.6	9.2	4.5	4.4	6.3	2.9	4.0	5.7	2.6	6.2	9.0	4.0
Malignant neoplasm of pancreas (C25)															
2000 .....	10.6	12.1	9.4	7.4	8.3	6.7	10.8	12.4	9.6	10.6	12.2	9.3	14.5	16.0	13.3
1999 .....	10.7	12.3	9.4	7.1	7.9	6.5	10.9	12.5	9.6	10.6	12.3	9.3	14.6	16.8	12.9
Malignant neoplasm of larynx (C32)															
2000 .....	1.4	2.5	0.5	0.8	1.7	*	1.4	2.6	0.6	1.3	2.4	0.5	2.9	5.6	1.0
1999 .....	1.4	2.6	0.5	1.1	2.2	0.3	1.4	2.6	0.6	1.3	2.3	0.5	2.7	5.2	0.9
Malignant neoplasms of trachea, bronchus and lung (C33-C34)															
2000 .....	56.5	76.6	41.8	22.0	34.0	13.1	58.7	79.2	43.7	58.9	78.1	44.8	66.1	102.5	41.4
1999 .....	56.0	77.0	40.8	22.5	35.2	13.1	58.0	79.4	42.5	58.0	78.2	43.4	67.0	104.8	41.7
Malignant melanoma of skin (C43)															
2000 .....	2.7	3.8	1.8	0.7	1.0	0.5	2.8	4.0	1.9	3.2	4.5	2.2	0.6	0.6	0.5
1999 .....	2.7	3.8	1.8	0.7	0.9	0.5	2.8	4.0	1.9	3.2	4.5	2.1	0.4	0.5	0.3
Malignant neoplasm of breast (C50)															
2000 .....	15.4	0.4	27.1	8.8	*	15.8	15.8	0.4	27.8	15.3	0.4	27.2	21.3	0.6	36.0
1999 .....	15.3	0.3	27.0	8.6	*	15.4	15.7	0.3	27.7	15.3	0.3	27.0	21.7	0.7	36.7
Malignant neoplasm of cervix uteri (C53)															
2000 .....	1.5	...	2.8	2.0	...	3.8	1.5	...	2.7	1.3	...	2.4	3.3	...	5.7
1999 .....	1.6	...	2.9	1.8	...	3.4	1.5	...	2.8	1.3	...	2.5	3.2	...	5.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)															
2000 .....	2.4	...	4.2	1.6	...	2.8	2.4	...	4.3	2.3	...	3.9	4.5	...	7.6
1999 .....	2.4	...	4.1	1.6	...	2.9	2.4	...	4.2	2.3	...	4.0	4.3	...	7.1

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Malignant neoplasm of ovary (C56)</b>															
2000 .....	5.1	...	9.0	3.3	...	5.8	5.2	...	9.2	5.4	...	9.6	4.4	...	7.5
1999 .....	5.0	...	8.9	2.8	...	5.1	5.2	...	9.1	5.3	...	9.5	4.5	...	7.6
<b>Malignant neoplasm of prostate (C61)</b>															
2000 .....	11.3	29.7	...	7.0	17.6	...	11.5	30.3	...	10.4	27.7	...	24.6	66.9	...
1999 .....	11.7	30.9	...	7.5	19.0	...	11.9	31.5	...	10.9	28.8	...	24.9	68.1	...
<b>Malignant neoplasms of kidney and renal pelvis (C64-C65)</b>															
2000 .....	4.3	6.1	2.9	3.2	4.7	2.0	4.3	6.2	2.9	4.4	6.3	2.9	4.3	6.3	2.9
1999 .....	4.1	5.9	2.7	3.3	4.7	2.1	4.1	6.0	2.8	4.2	6.0	2.8	4.3	6.2	3.0
<b>Malignant neoplasm of bladder (C67)</b>															
2000 .....	4.3	7.5	2.3	2.0	3.4	1.0	4.5	7.7	2.4	4.7	8.1	2.4	3.8	5.5	2.8
1999 .....	4.4	7.6	2.3	2.0	3.5	0.9	4.5	7.8	2.4	4.6	8.1	2.4	4.1	5.7	3.1
<b>Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)</b>															
2000 .....	4.6	5.6	3.7	2.9	3.4	2.4	4.7	5.8	3.8	5.1	6.2	4.1	2.8	3.4	2.4
1999 .....	4.7	5.7	3.9	2.8	3.0	2.5	4.8	5.9	4.0	5.2	6.3	4.3	2.8	3.4	2.4
<b>Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)</b>															
2000 .....	20.5	25.8	16.7	14.1	17.3	11.6	20.9	26.2	16.9	21.1	26.7	17.1	20.6	25.4	17.4
1999 .....	20.7	26.1	16.9	14.1	16.2	12.4	21.1	26.6	17.1	21.3	26.9	17.2	21.0	26.1	17.6
<b>Hodgkin's disease (C81)</b>															
2000 .....	0.5	0.6	0.4	0.5	0.7	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.4	0.3
1999 .....	0.5	0.7	0.4	0.6	0.8	0.5	0.5	0.6	0.4	0.6	0.7	0.4	0.4	0.6	0.3
<b>Non-Hodgkin's lymphoma (C82-C85)</b>															
2000 .....	8.3	10.2	6.8	5.7	6.9	4.8	8.4	10.3	6.9	8.7	10.8	7.3	5.4	6.8	4.4
1999 .....	8.4	10.3	7.0	5.7	6.5	5.0	8.6	10.5	7.1	8.9	10.9	7.4	5.8	7.6	4.6
<b>Leukemia (C91-C95)</b>															
2000 .....	7.8	10.2	6.0	5.0	6.3	4.0	7.9	10.4	6.1	8.0	10.6	6.2	6.9	9.1	5.5
1999 .....	7.7	10.2	6.0	4.9	5.8	4.2	7.8	10.4	6.1	8.0	10.6	6.2	6.9	8.9	5.6
<b>Multiple myeloma and immunoproliferative neoplasms (C88,C90)</b>															
2000 .....	4.0	4.9	3.5	2.9	3.3	2.5	4.1	4.9	3.5	3.8	4.6	3.2	7.9	9.0	7.1
1999 .....	4.1	4.9	3.4	2.9	3.1	2.7	4.1	5.0	3.5	3.8	4.7	3.1	7.8	9.0	7.0

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)</b>															
2000 .....	0.0	0.0	0.0	*	*	*	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
1999 .....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
<b>All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)</b>															
2000 .....	22.9	27.3	19.7	15.5	17.8	13.8	23.3	27.9	20.0	23.1	27.6	19.8	27.3	33.4	23.0
1999 .....	24.4	29.3	20.8	16.2	18.7	14.4	24.8	29.9	21.2	24.6	29.6	20.9	29.5	35.9	25.0
<b>In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)</b>															
2000 .....	4.9	5.9	4.2	3.0	3.3	2.9	5.0	6.1	4.3	5.1	6.2	4.4	4.7	5.4	4.3
1999 .....	4.9	6.0	4.1	2.9	3.0	2.8	5.0	6.2	4.2	5.0	6.3	4.2	5.0	5.4	4.6
<b>Anemias (D50-D64)</b>															
2000 .....	1.6	1.5	1.7	0.8	0.8	0.9	1.7	1.6	1.7	1.4	1.4	1.4	3.6	3.5	3.7
1999 .....	1.6	1.7	1.6	1.0	1.2	0.9	1.7	1.7	1.7	1.4	1.4	1.4	3.8	3.8	3.7
<b>Diabetes mellitus (E10-E14)</b>															
2000 .....	25.2	27.6	23.3	32.4	34.6	30.5	24.7	27.0	22.8	22.0	25.0	19.7	51.1	49.2	51.7
1999 .....	25.2	27.7	23.3	33.6	34.5	32.6	24.7	27.2	22.7	21.9	25.0	19.5	51.4	50.1	51.7
<b>Nutritional deficiencies (E40-E64)</b>															
2000 .....	1.5	1.4	1.6	1.0	1.0	1.0	1.5	1.4	1.6	1.5	1.3	1.6	2.5	2.6	2.3
1999 .....	1.6	1.5	1.6	1.0	1.0	1.0	1.6	1.5	1.6	1.5	1.4	1.6	2.8	3.1	2.5
<b>Malnutrition (E40-E46)</b>															
2000 .....	1.4	1.3	1.5	0.9	0.9	0.9	1.4	1.3	1.5	1.4	1.2	1.4	2.3	2.5	2.2
1999 .....	1.5	1.4	1.5	1.0	1.0	1.0	1.5	1.4	1.5	1.4	1.3	1.5	2.5	2.9	2.3
<b>Other nutritional deficiencies (E50-E64)</b>															
2000 .....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999 .....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
<b>Meningitis (G00,G03)</b>															
2000 .....	0.3	0.3	0.2	0.2	0.2	0.1	0.3	0.3	0.2	0.2	0.3	0.2	0.7	0.8	0.6
1999 .....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6
<b>Parkinson's disease (G20-G21)</b>															
2000 .....	5.7	8.6	4.0	3.0	4.3	2.2	5.8	8.8	4.1	6.2	9.3	4.3	2.6	4.1	1.7
1999 .....	5.4	8.2	3.7	2.9	4.2	2.1	5.5	8.4	3.8	5.8	9.0	4.0	2.4	3.7	1.8

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Alzheimer's disease (G30)</b>															
2000 .....	18.0	14.7	19.5	8.2	7.1	8.8	18.4	15.2	20.0	19.2	15.8	20.8	13.4	11.1	14.5
1999 .....	16.5	14.0	17.6	7.7	6.4	8.4	16.9	14.4	18.0	17.6	15.1	18.8	11.6	9.9	12.4
<b>Major cardiovascular diseases (I00-I78)</b>															
2000 .....	339.9	400.5	291.3	217.9	258.2	185.1	345.8	407.8	296.2	337.5	400.5	286.8	452.2	517.7	402.1
1999 .....	350.9	414.8	300.3	229.2	273.0	194.1	356.5	421.6	304.9	348.2	414.0	295.8	461.4	533.5	407.2
<b>Diseases of heart (I00-I09,I11,I13, I20-I51)</b>															
2000 .....	257.9	315.0	213.0	165.4	200.5	137.1	262.4	321.0	216.6	257.5	317.2	210.4	334.6	392.7	291.0
1999 .....	267.8	328.1	220.9	176.1	212.7	146.5	272.0	333.6	224.2	266.9	329.5	218.1	344.3	408.2	297.0
<b>Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)</b>															
2000 .....	1.3	1.0	1.6	0.8	0.6	1.0	1.3	1.0	1.6	1.3	1.0	1.6	1.0	0.7	1.3
1999 .....	1.4	0.9	1.6	1.0	0.4	1.4	1.4	0.9	1.6	1.4	1.0	1.7	1.1	0.8	1.3
<b>Hypertensive heart disease (I11)</b>															
2000 .....	8.6	8.6	8.1	6.0	6.5	5.4	8.7	8.7	8.3	7.1	6.9	6.8	24.5	26.6	22.2
1999 .....	8.4	8.3	8.0	6.0	6.3	5.6	8.5	8.4	8.1	6.9	6.7	6.7	23.7	26.1	21.4
<b>Hypertensive heart and renal disease (I13)</b>															
2000 .....	1.0	1.1	1.0	0.9	1.0	0.8	1.0	1.1	1.0	0.8	0.8	0.7	3.4	3.7	3.2
1999 .....	1.2	1.3	1.1	1.0	1.0	0.9	1.2	1.3	1.2	0.9	1.0	0.8	4.3	4.5	4.1
<b>Ischemic heart diseases (I20-I25)</b>															
2000 .....	186.9	237.7	147.9	128.8	159.7	104.3	189.7	241.6	149.9	188.1	241.6	146.8	224.3	268.8	191.7
1999 .....	195.6	249.0	154.9	138.4	170.4	112.8	198.1	252.7	156.6	196.6	252.8	153.7	231.1	278.4	196.6
<b>Acute myocardial infarction (I21-I22)</b>															
2000 .....	70.0	90.1	54.7	45.8	57.1	37.0	71.3	92.0	55.5	71.0	92.5	54.3	83.5	99.0	72.4
1999 .....	73.6	95.4	57.1	49.2	61.0	39.8	74.9	97.3	57.9	74.7	98.1	56.8	85.8	102.6	73.9
<b>Other acute ischemic heart diseases (I24)</b>															
2000 .....	1.2	1.5	0.9	0.4	0.6	0.3	1.3	1.6	1.0	1.2	1.5	0.9	1.9	2.5	1.4
1999 .....	1.2	1.5	0.9	0.3	0.4	0.3	1.2	1.6	1.0	1.2	1.5	0.9	1.8	2.5	1.3
<b>Other forms of chronic ischemic heart disease (I20,I25)</b>															
2000 .....	115.7	146.1	92.3	82.6	102.0	66.9	117.1	148.0	93.4	115.9	147.6	91.6	139.0	167.3	117.9
1999 .....	120.8	152.1	96.9	88.9	109.1	72.7	122.0	153.8	97.7	120.7	153.2	96.0	143.5	173.3	121.3
<b>Atherosclerotic cardiovascular disease, so described (I25.0)</b>															
2000 .....	25.2	31.9	19.6	18.3	25.0	12.9	25.4	32.1	19.9	23.7	29.8	18.5	44.7	58.9	34.4
1999 .....	26.5	33.3	20.9	20.2	27.2	14.5	26.7	33.4	21.1	24.8	30.9	19.7	47.0	61.5	36.3

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)															
2000 .....	90.5	114.2	72.8	64.3	77.1	54.0	91.6	116.0	73.5	92.2	117.8	73.1	94.2	108.4	83.5
1999 .....	94.2	118.7	75.9	68.7	81.8	58.2	95.2	120.4	76.6	95.9	122.2	76.3	96.5	111.8	85.0
Other heart diseases (I26-I51)															
2000 .....	60.0	66.7	54.4	28.8	32.7	25.7	61.7	68.6	55.8	60.1	66.9	54.4	81.3	92.9	72.6
1999 .....	61.2	68.5	55.2	29.8	34.6	25.8	62.8	70.3	56.6	61.1	68.2	55.1	84.1	98.4	73.7
Acute and subacute endocarditis (I33)															
2000 .....	0.4	0.5	0.3	0.3	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.7
1999 .....	0.4	0.5	0.3	0.3	0.5	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.0	0.6
Diseases of pericardium and acute myocarditis (I30-I31,I40)															
2000 .....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.2	0.3	0.3	0.2	0.4	0.6	0.4
1999 .....	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.6	0.6	0.5
Heart failure (I50)															
2000 .....	20.2	20.9	19.4	8.8	9.2	8.4	20.8	21.6	19.9	20.8	21.7	19.9	22.7	23.9	21.8
1999 .....	20.3	21.0	19.5	9.2	9.5	8.8	20.8	21.6	20.0	20.8	21.6	20.0	23.2	25.0	21.9
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)															
2000 .....	39.1	44.9	34.4	19.4	22.8	16.7	40.2	46.2	35.4	38.7	44.4	34.0	57.3	67.5	49.7
1999 .....	40.2	46.5	35.1	20.0	24.4	16.5	41.3	47.8	36.0	39.6	45.8	34.6	59.5	71.9	50.6
Essential (primary) hypertension and hypertensive renal disease (I10,I12)															
2000 .....	6.6	6.4	6.5	5.3	5.7	4.9	6.6	6.4	6.6	5.6	5.3	5.6	16.9	17.2	16.4
1999 .....	6.3	6.1	6.2	5.2	5.5	4.9	6.3	6.1	6.3	5.2	5.1	5.2	16.6	17.4	15.8
Cerebrovascular diseases (I60-I69)															
2000 .....	60.8	61.1	59.7	39.2	42.4	36.4	61.8	61.9	60.7	59.4	59.2	58.6	84.5	89.5	80.1
1999 .....	61.8	62.4	60.5	40.0	44.6	36.3	62.8	63.1	61.6	60.5	60.5	59.6	84.5	89.7	80.0
Atherosclerosis (I70)															
2000 .....	5.2	5.3	5.0	2.9	3.0	2.8	5.3	5.5	5.1	5.4	5.6	5.2	4.9	5.3	4.7
1999 .....	5.5	5.6	5.4	3.1	3.6	2.8	5.6	5.7	5.5	5.8	5.8	5.6	5.1	5.7	4.7
Other diseases of circulatory system (I71-I78)															
2000 .....	9.4	12.7	7.1	5.0	6.5	3.9	9.7	13.1	7.3	9.6	13.2	7.1	11.3	13.0	9.9
1999 .....	9.5	12.7	7.2	4.9	6.7	3.6	9.7	13.0	7.4	9.7	13.2	7.2	10.8	12.4	9.7
Aortic aneurysm and dissection (I71)															
2000 .....	5.7	8.6	3.7	2.8	4.1	1.8	5.9	8.9	3.8	6.0	9.1	3.7	5.4	6.8	4.3
1999 .....	5.8	8.7	3.8	2.5	3.9	1.5	6.0	9.0	3.9	6.1	9.3	3.9	5.1	6.5	4.1

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)															
2000 .....	3.7	4.1	3.4	2.2	2.4	2.0	3.8	4.2	3.5	3.6	4.1	3.3	5.9	6.2	5.6
1999 .....	3.7	4.0	3.5	2.4	2.8	2.1	3.7	4.0	3.5	3.6	3.9	3.4	5.8	6.0	5.6
Other disorders of circulatory system (I80-I99)															
2000 .....	1.7	1.7	1.6	1.2	1.4	1.1	1.7	1.7	1.6	1.6	1.6	1.5	3.0	3.1	2.9
1999 .....	1.5	1.5	1.4	1.1	1.1	1.1	1.5	1.5	1.5	1.4	1.4	1.3	2.5	2.7	2.3
Influenza and pneumonia (J10-J18)															
2000 .....	23.7	28.1	20.9	17.0	20.7	14.4	23.9	28.4	21.2	23.7	27.9	21.1	26.4	33.5	21.8
1999 .....	23.6	28.0	20.8	15.6	18.6	13.5	23.9	28.4	21.1	23.7	28.0	21.1	26.1	33.0	21.7
Influenza (J10-J11)															
2000 .....	0.6	0.6	0.6	0.3	0.3	0.2	0.6	0.7	0.6	0.7	0.7	0.7	0.4	0.4	0.3
1999 .....	0.6	0.6	0.6	0.3	0.2	0.3	0.6	0.6	0.6	0.6	0.7	0.6	0.3	0.3	0.3
Pneumonia (J12-J18)															
2000 .....	23.0	27.5	20.2	16.7	20.4	14.1	23.3	27.7	20.5	23.0	27.2	20.4	26.0	33.1	21.5
1999 .....	22.9	27.4	20.2	15.3	18.3	13.3	23.2	27.7	20.5	23.0	27.3	20.4	25.8	32.8	21.4
Other acute lower respiratory infections (J20-J22)															
2000 .....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2
1999 .....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Acute bronchitis and bronchiolitis (J20-J21)															
2000 .....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999 .....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Unspecified acute lower respiratory infection (J22)															
2000 .....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999 .....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Chronic lower respiratory diseases (J40-J47)															
2000 .....	44.3	54.9	37.9	17.7	23.6	13.8	45.8	56.6	39.2	47.7	58.1	41.5	32.5	47.1	23.8
1999 .....	45.8	58.1	38.2	19.9	27.3	15.3	47.1	59.8	39.4	48.9	61.3	41.5	34.5	51.4	24.5
Bronchitis, chronic and unspecified (J40-J42)															
2000 .....	0.4	0.5	0.4	0.3	0.4	0.3	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.3	0.3
1999 .....	0.4	0.5	0.4	0.3	*	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
Emphysema (J43)															
2000 .....	6.1	7.7	5.0	1.8	2.7	1.2	6.3	8.0	5.2	6.7	8.3	5.6	3.9	6.1	2.5
1999 .....	6.5	8.6	5.2	2.1	3.2	1.3	6.8	8.9	5.5	7.2	9.2	5.9	4.1	6.8	2.4

See footnotes at end of table.



**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Asthma (J45-J46)</b>															
2000 .....	1.6	1.3	1.8	1.5	1.3	1.6	1.6	1.3	1.9	1.3	1.0	1.5	4.1	3.6	4.3
1999 .....	1.7	1.4	2.0	1.7	1.2	2.0	1.7	1.4	2.0	1.4	1.0	1.7	4.1	3.7	4.3
<b>Other chronic lower respiratory diseases (J44,J47)</b>															
2000 .....	36.2	45.4	30.6	14.2	19.2	10.9	37.4	46.8	31.8	39.3	48.3	33.9	24.3	37.0	16.7
1999 .....	37.0	47.8	30.6	15.9	22.5	11.6	38.1	49.0	31.6	39.9	50.6	33.6	26.0	40.5	17.4
<b>Pneumoconioses and chemical effects (J60-J66,J68)</b>															
2000 .....	0.4	1.1	0.0	0.2	0.4	*	0.4	1.1	0.0	0.5	1.2	0.0	0.3	0.7	*
1999 .....	0.4	1.0	0.0	*	*	*	0.4	1.1	0.0	0.4	1.1	0.0	0.3	0.8	*
<b>Pneumonitis due to solids and liquids (J69)</b>															
2000 .....	6.0	8.3	4.7	3.3	4.3	2.7	6.2	8.5	4.8	6.1	8.6	4.8	6.9	9.3	5.5
1999 .....	5.6	7.9	4.4	3.0	3.7	2.6	5.8	8.1	4.4	5.7	8.1	4.4	7.0	9.7	5.4
<b>Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)</b>															
2000 .....	9.3	11.1	8.0	7.2	8.2	6.4	9.4	11.3	8.1	9.4	11.3	8.1	9.6	11.3	8.4
1999 .....	9.2	11.1	8.0	7.3	8.6	6.3	9.3	11.2	8.0	9.3	11.2	7.9	9.8	11.3	8.8
<b>Peptic ulcer (K25-K28)</b>															
2000 .....	1.6	2.0	1.4	1.2	1.3	1.0	1.7	2.0	1.4	1.6	2.0	1.4	1.8	2.4	1.4
1999 .....	1.7	2.0	1.4	1.1	1.6	0.8	1.7	2.0	1.5	1.7	2.0	1.5	1.9	2.4	1.5
<b>Diseases of appendix (K35-K38)</b>															
2000 .....	0.2	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999 .....	0.1	0.2	0.1	0.1	0.2	*	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.1
<b>Hernia (K40-K46)</b>															
2000 .....	0.6	0.5	0.6	0.5	0.4	0.5	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.6	0.6
1999 .....	0.5	0.5	0.5	0.4	0.4	0.5	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>															
2000 .....	9.6	13.5	6.2	15.7	23.7	8.6	9.1	12.6	6.1	9.0	12.5	5.9	9.8	14.2	6.5
1999 .....	9.7	13.7	6.1	15.3	23.0	8.5	9.2	12.9	6.0	9.1	12.7	5.8	10.5	15.6	6.6
<b>Alcoholic liver disease (K70)</b>															
2000 .....	4.4	6.9	2.2	7.6	14.1	2.0	4.1	6.3	2.2	4.0	6.2	2.0	4.9	7.4	2.9
1999 .....	4.4	7.0	2.1	7.4	13.3	2.3	4.2	6.5	2.1	4.0	6.2	2.0	5.3	8.5	2.9
<b>Other chronic liver disease and cirrhosis (K73-K74)</b>															
2000 .....	5.3	6.6	4.0	8.0	9.6	6.6	5.0	6.4	3.9	5.0	6.4	3.9	5.0	6.8	3.6
1999 .....	5.3	6.7	4.0	7.9	9.7	6.2	5.1	6.5	3.9	5.1	6.4	3.9	5.2	7.1	3.7

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Cholelithiasis and other disorders of gallbladder (K80-K82)</b>															
2000 .....	1.0	1.2	0.9	1.2	1.4	0.9	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.0	1.3
1999 .....	1.0	1.2	1.0	1.3	1.4	1.2	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
<b>Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19, N25-N27)</b>															
2000 .....	13.5	16.6	11.6	10.1	11.8	9.0	13.7	16.8	11.7	12.1	15.4	10.1	29.6	33.2	27.3
1999 .....	13.1	16.2	11.2	10.4	12.9	8.8	13.2	16.4	11.3	11.6	14.8	9.7	29.4	33.8	26.6
<b>Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)</b>															
2000 .....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
1999 .....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	*
<b>Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07, N26)</b>															
2000 .....	0.2	0.3	0.2	0.1	*	*	0.2	0.3	0.2	0.2	0.2	0.1	0.4	0.4	0.3
1999 .....	0.2	0.3	0.2	0.2	*	*	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.3
<b>Renal failure (N17-N19)</b>															
2000 .....	13.2	16.3	11.4	9.9	11.4	8.9	13.4	16.5	11.5	11.8	15.1	9.9	29.1	32.6	26.8
1999 .....	12.8	15.9	11.0	10.1	12.6	8.5	12.9	16.0	11.1	11.3	14.4	9.5	28.8	33.2	26.1
<b>Other disorders of kidney (N25,N27)</b>															
2000 .....	0.0	0.0	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
1999 .....	0.0	*	0.0	*	*	*	0.0	*	0.0	*	*	*	*	*	*
<b>Infections of kidney (N10-N12,N13.6, N15.1)</b>															
2000 .....	0.3	0.2	0.4	0.2	*	0.3	0.3	0.2	0.4	0.3	0.2	0.3	0.4	0.3	0.5
1999 .....	0.3	0.2	0.3	0.3	0.3	0.4	0.3	0.2	0.3	0.3	0.2	0.4	0.4	0.3	0.4
<b>Hyperplasia of prostate (N40)</b>															
2000 .....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999 .....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
<b>Inflammatory diseases of female pelvic organs (N70-N76)</b>															
2000 .....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999 .....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Pregnancy, childbirth and the puerperium (O00-O99)</b>															
2000 .....	0.1	...	0.3	0.2	...	0.5	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999 .....	0.2	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
<b>Pregnancy with abortive outcome (O00-O07)</b>															
2000 .....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999 .....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
<b>Other complications of pregnancy, childbirth and the puerperium (O10-O99)</b>															
2000 .....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.2	0.1	...	0.2	0.3	...	0.6
1999 .....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.2	0.1	...	0.1	0.4	...	0.8
<b>Certain conditions originating in the perinatal period (P00-P96)</b>															
2000 .....	5.0	5.6	4.5	4.0	4.4	3.6	5.2	5.8	4.6	3.7	4.1	3.2	12.4	13.8	11.0
1999 .....	5.1	5.7	4.6	4.1	4.5	3.8	5.3	5.8	4.7	3.7	4.1	3.3	13.1	14.5	11.6
<b>Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)</b>															
2000 .....	3.8	4.1	3.6	3.7	4.0	3.4	3.8	4.0	3.6	3.6	3.9	3.4	4.8	5.1	4.5
1999 .....	3.8	4.0	3.6	3.5	3.5	3.5	3.8	4.0	3.6	3.6	3.8	3.4	4.8	5.1	4.5
<b>Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)</b>															
2000 .....	11.5	13.1	9.8	7.5	10.0	5.1	11.7	13.1	10.1	11.0	12.4	9.5	17.2	20.3	14.5
1999 .....	9.8	10.9	8.5	4.9	6.0	3.9	10.1	11.3	8.8	9.4	10.3	8.2	16.2	19.8	13.2
<b>All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)</b>															
2000 .....	64.7	64.5	63.0	40.1	43.0	37.0	65.9	65.7	64.4	64.2	63.5	62.9	87.0	93.4	81.2
1999 .....	63.4	63.7	61.4	42.2	45.2	38.9	64.4	64.5	62.5	62.7	62.3	61.1	84.1	91.2	78.3
<b>Accidents (unintentional injuries) (V01-X59, Y85-Y86)</b>															
2000 .....	35.5	50.0	22.3	30.6	46.2	15.5	35.6	49.7	22.8	35.4	49.0	22.9	39.7	60.7	22.6
1999 .....	35.9	50.6	22.7	31.1	47.2	15.6	36.0	50.3	23.2	35.5	49.1	23.1	42.2	64.2	24.4
<b>Transport accidents (V01-V99, Y85)</b>															
2000 .....	16.9	24.2	10.2	16.6	24.9	8.5	16.9	23.9	10.3	16.9	23.7	10.5	17.6	27.6	9.3

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Transport accidents (V01-V99,Y85)															
1999 .....	17.0	24.1	10.4	16.2	24.1	8.5	17.0	23.9	10.6	16.9	23.7	10.7	18.1	28.0	10.0
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)															
2000 .....	15.7	22.2	9.7	15.5	23.1	8.1	15.6	21.9	9.9	15.6	21.6	10.0	16.6	25.7	9.0
1999 .....	15.5	21.8	9.8	15.0	21.9	8.1	15.5	21.6	9.9	15.4	21.3	10.0	16.8	25.6	9.5
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)															
2000 .....	0.5	0.8	0.3	0.7	1.2	0.2	0.5	0.8	0.3	0.5	0.8	0.2	0.6	1.1	0.3
1999 .....	0.8	1.2	0.4	0.9	1.6	0.3	0.7	1.2	0.4	0.8	1.1	0.4	0.8	1.3	0.3
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)															
2000 .....	0.7	1.2	0.2	0.4	0.7	*	0.7	1.3	0.2	0.8	1.3	0.3	0.4	0.8	*
1999 .....	0.7	1.2	0.3	0.3	0.5	*	0.7	1.2	0.3	0.8	1.3	0.3	0.6	1.0	0.2
Nontransport accidents (W00-X59,Y86)															
2000 .....	18.5	25.8	12.1	14.0	21.2	7.0	18.7	25.9	12.5	18.5	25.3	12.4	22.0	33.2	13.2
1999 .....	18.9	26.4	12.3	14.9	23.2	7.0	19.0	26.3	12.6	18.5	25.5	12.4	24.0	36.2	14.4
Falls (W00-W19)															
2000 .....	4.8	6.5	3.6	3.7	5.3	2.3	4.9	6.5	3.7	5.1	6.7	3.8	3.0	4.6	2.0
1999 .....	4.8	6.6	3.6	3.6	5.4	2.2	4.9	6.6	3.6	5.0	6.7	3.8	3.5	5.3	2.2
Accidental discharge of firearms (W32-W34)															
2000 .....	0.3	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1
1999 .....	0.3	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.8	0.1
Accidental drowning and submersion (W65-W74)															
2000 .....	1.3	2.0	0.5	1.2	1.9	0.4	1.3	2.0	0.5	1.1	1.8	0.5	1.6	2.8	0.6
1999 .....	1.3	2.1	0.5	1.3	2.2	0.4	1.3	2.0	0.5	1.2	1.8	0.5	1.8	3.1	0.6
Accidental exposure to smoke, fire and flames (X00-X09)															
2000 .....	1.2	1.6	0.9	0.7	0.9	0.5	1.3	1.7	1.0	1.1	1.4	0.8	3.2	4.3	2.4
1999 .....	1.2	1.6	0.9	0.7	1.0	0.4	1.3	1.7	0.9	1.1	1.4	0.8	3.2	4.2	2.4

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental poisoning and exposure to noxious substances (X40-X49)															
2000 .....	4.6	6.7	2.6	4.4	7.2	1.5	4.6	6.6	2.6	4.6	6.5	2.6	6.3	9.8	3.4
1999 .....	4.5	6.6	2.3	5.2	8.7	1.7	4.3	6.3	2.4	4.2	6.0	2.3	6.5	10.3	3.3
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)															
2000 .....	6.3	8.4	4.5	3.9	5.7	2.2	6.4	8.6	4.6	6.4	8.4	4.6	7.4	11.0	4.8
1999 .....	6.8	9.0	4.8	3.8	5.5	2.4	6.9	9.2	5.0	6.8	9.1	4.9	8.6	12.5	5.7
Intentional self-harm (suicide) (X60-X84,Y87.0)															
2000 .....	10.6	18.1	4.0	6.1	10.8	1.8	11.1	18.8	4.2	12.1	20.2	4.7	5.8	10.5	1.9
1999 .....	10.7	18.2	4.1	6.1	10.7	1.9	11.1	18.9	4.3	12.1	20.3	4.7	5.9	10.8	1.7
Intentional self-harm (suicide) by discharge of firearms (X72-X74)															
2000 .....	6.0	11.2	1.5	2.8	5.3	0.5	6.3	11.8	1.6	6.9	12.8	1.8	3.3	6.3	0.7
1999 .....	6.1	11.4	1.5	2.9	5.4	0.6	6.4	11.9	1.6	7.0	12.9	1.8	3.4	6.9	0.6
Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)															
2000 .....	4.6	6.9	2.5	3.3	5.5	1.3	4.8	7.0	2.7	5.1	7.4	2.9	2.6	4.2	1.1
1999 .....	4.6	6.8	2.6	3.2	5.2	1.3	4.8	7.0	2.7	5.1	7.4	2.9	2.5	4.0	1.1
Assault (homicide) (X85-Y09,Y87.1)															
2000 .....	6.1	9.3	2.8	8.4	13.6	3.0	5.7	8.6	2.8	2.8	3.6	1.9	21.8	37.6	7.5
1999 .....	6.2	9.4	2.9	8.4	13.8	2.8	5.8	8.6	2.9	2.9	3.9	2.0	21.4	36.4	7.8
Assault (homicide) by discharge of firearms (X93-X95)															
2000 .....	3.9	6.5	1.3	5.4	9.3	1.3	3.6	6.0	1.3	1.5	2.0	0.9	15.5	28.6	3.3
1999 .....	4.0	6.5	1.4	5.5	9.5	1.4	3.7	6.0	1.3	1.5	2.2	0.9	15.3	27.8	3.7
Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)															
2000 .....	2.1	2.8	1.5	2.9	4.2	1.6	2.0	2.6	1.5	1.3	1.6	1.0	6.3	8.9	4.2
1999 .....	2.2	2.9	1.6	2.8	4.3	1.4	2.1	2.6	1.6	1.4	1.7	1.1	6.1	8.6	4.1
Legal intervention (Y35,Y89.0)															
2000 .....	0.1	0.3	*	0.1	0.3	*	0.1	0.2	*	0.1	0.2	*	0.3	0.7	*
1999 .....	0.2	0.3	*	0.2	0.4	*	0.1	0.3	*	0.1	0.2	*	0.4	0.7	*

See footnotes at end of table.

**Orig293. Age-Adjusted Death Rates for 113 Selected Causes, by Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2000 - Con.**

[Age-adjusted death rates per 100,000 U.S. standard population based on year 2000 standard in specified group. Populations used for computing death rates are postcensal estimates based on the 1990 census, estimated as of July 1. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All origins <sup>1</sup>			Hispanic			Non-Hispanic <sup>2</sup>			Non-Hispanic, white			Non-Hispanic, black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)</b>															
2000 .....	1.4	1.9	0.8	0.7	1.0	0.3	1.5	2.0	0.9	1.4	1.9	0.9	2.0	3.2	1.0
1999 .....	1.4	2.0	0.9	0.8	1.3	0.3	1.5	2.1	0.9	1.4	1.9	0.9	2.1	3.5	1.0
<b>Discharge of firearms, undetermined intent (Y22-Y24)</b>															
2000 .....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999 .....	0.1	0.2	0.0	0.1	0.2	*	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*
<b>Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)</b>															
2000 .....	1.3	1.8	0.8	0.6	0.9	0.3	1.4	1.9	0.9	1.3	1.8	0.9	1.9	2.9	1.0
1999 .....	1.3	1.8	0.9	0.7	1.1	0.3	1.4	1.8	0.9	1.3	1.7	0.9	1.9	3.1	0.9
<b>Operations of war and their sequelae (Y36,Y89.1)</b>															
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
<b>Complications of medical and surgical care (Y40-Y84,Y88)</b>															
2000 .....	1.1	1.2	1.0	0.7	0.8	0.6	1.1	1.3	1.0	1.1	1.2	1.0	2.0	1.9	2.0
1999 .....	1.0	1.2	1.0	0.5	0.6	0.4	1.1	1.2	1.0	1.0	1.1	0.9	1.8	2.0	1.7

0.0 Quantity more than zero but less than 0.05.  
 Figure does not meet standards of reliability or precision.

1 . . . Category not applicable.

<sup>1</sup> All origins include origin not stated; specified origins exclude origin not stated.

<sup>2</sup> Includes races other than white and black.

Source: CDC/NCHS, National Vital Statistics System, Mortality.